



FOR 24 HOUR  
SPECIALIST  
ADVICE IN  
EMERGENCY  
ONLY  
**PHONE**  
**1 800 033 111**

**Fly Bait**

**DIRECTIONS FOR USE**  
**RESTRAINTS: DO NOT** apply in a manner which may cause contamination of animal feed.  
**DO NOT** apply spray treatment outdoors if rain is expected within 48 hours.  
**DO NOT** apply onto material which birds are likely to feed on directly (i.e. edible refuse).

SITUATION	PEST	RATE	CRITICAL COMMENTS
This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.			
<b>OUTSIDE:</b> Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas <b>AROUND:</b> Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children) <b>INSIDE</b> Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)	Nuisance flies including: Adult houseflies ( <i>Musca domestica</i> ) and lesser houseflies ( <i>Fannia canicularis</i> )	250g / 2L water (spray on application)	Mix 250g of QuickBayt® Spray with 2L of water and apply as a coarse spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.  For floor areas of 100m <sup>2</sup> it is recommended that 2L of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor area greater than 100m <sup>2</sup> needs to be recalculated on the basis of 2L mixed bait per 100m <sup>2</sup> floor area.  <b>Note: Around feedlot manure heaps DO NOT</b> apply bait on manure heaps or dispose of treated targets with manure. <b>See also Note on Application below.</b>
		250g / 200mL water (paint on application)	To prepare a paste for painting, mix 250g of QuickBayt® Spray with 200mL of water. The mix can be applied with a brush to spots where flies rest (e.g. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water. For floor areas of 100m <sup>2</sup> it is recommended that 200mL of mix is applied in 50 to 100 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 100m <sup>2</sup> need to be recalculated on the basis of 200mL mixed bait per 100m <sup>2</sup> floor area.  <b>See also Note on Application below.</b>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS**  
QuickBayt® Spray can be applied in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. For best results time applications to begin at the start of the season before fly populations have reached their peak.  
  
Fly control should be considered part of an integrated strategy which incorporates good sanitation, appropriate management of breeding sites (e.g. manure and other decaying organic matter) and exclusion measures for preventing access of flies into internal areas.

**Note on Application**  
Avoid applying this product to dusty/dirty surfaces. If the treated area becomes covered by dust/dirt, the product can be reactivated by gently spraying water mist on the treated surfaces.

**DO NOT** store diluted insecticide. Prepare a fresh solution as necessary.  
**DO NOT** spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

**Re-baiting:**  
Re-apply bait when flies are no longer visibly affected in areas where the product has been applied (usually after about 4 to 6 weeks). For season-long control, monthly applications may be required, depending on the situation and duration of the fly season.

**Cleaning Spray Equipment**  
Clean spray equipment by rinsing thoroughly with lukewarm water.  
**DO NOT** allow washings to enter sewer drains or nearby water bodies.

**WARNING:**  
In some situations mould may develop (some time after application) on areas where this product has been used. If appearance of mould on a wall or fixture is a concern in the area of application then it is recommended that a surrogate target such as a disposable piece of timber, card or other surface is treated in its place. A suitable detergent and cloth may also be used to remove mould in some areas. Dispose of cloth after use.

**Insecticide Resistance Warning**  
**GROUP 4A INSECTICIDE**

For insecticide resistance management, QuickBayt® Spray is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to QuickBayt® Spray and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if QuickBayt® Spray or other Group 4A insecticides are used repeatedly. The effectiveness of QuickBayt® Spray on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of QuickBayt® Spray to control resistant insects. For further information on managing insecticide resistance contact your local supplier or Bayer Animal Health representative.

**PRECAUTIONS**  
**DO NOT** allow contact with raw or processed food, utensils or packaging.  
**DO NOT** apply to surfaces within the reach of animals and children.  
**DO NOT** spray directly on animals, feed or drinking supplies, or on milking equipment. **DO NOT** apply past the point of run-off.

**PROTECTION OF LIVESTOCK**  
**DO NOT** apply bait in locations accessible to domestic animals or livestock. Imidacloprid is dangerous to bees. QuickBayt® Spray will kill bees in hives which are over-sprayed or reached with spray drift.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT**  
**DO NOT** apply bait in locations accessible to birds.  
**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**  
Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Single-rinse or shake remainder into spray tank/water. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on site. Do not burn empty containers or product.

**SAFETY DIRECTIONS**  
When opening the container, preparing and using the prepared spray or paste wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length chemical resistant gloves. Wash hands after use.

**FIRST AID**  
If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

**SAFETY DATA SHEET**  
Additional information is listed in the Safety Data Sheet available from [www.growsolutions.com.au](http://www.growsolutions.com.au)

**EXCLUSION OF LIABILITY**  
This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

QuickBayt® is a registered trade mark of the Bayer Group.  
APVMA Approval No. 61737/119534



Distributed by:  
**Bayer**

Bayer Australia Ltd  
ACN 000 138 714  
875 Pacific Highway  
Pymble NSW 2073

Technical enquiries: 1800 678 368  
[www.growsolutions.com.au](http://www.growsolutions.com.au)

AU80001738B 05/19

Batch:  
DOM:  
Expiry:

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Fly Bait**

**ACTIVE CONSTITUENTS:**  
100g/kg IMIDACLOPRID, 0.84g/kg (Z)-9 TRICOSENE

**GROUP 4A INSECTICIDE**

A water dispersible granule formulation to be mixed with water and used as a spray bait formulation for the control of adult houseflies and lesser houseflies (including organophosphate resistant strains) in commercial, industrial and domestic areas, as per the DIRECTIONS FOR USE. Will visibly kill flies for 4 weeks.



**1kg Net**

Contains 4 x 250g measure packs which it is illegal to sell separately.

APVMA Approval No. 61737/119534